

Effect of basketball specific endurance circuit training on aerobic capacity and heart rate of high school male basketball players

B. Chittibabu^{a,*} and N. Akilan^a

^aAssistant professor, Department of Physical Education and Sports Sciences, Annamalai University, Chidambaram, Tamil Nadu-608002, India

**Corresponding Author Ph: 09443531508; Email: b.chitti@hotmail.com*

DOI: 10.26524/1345

Abstract: The purpose of the study was to evaluate the effectiveness of a basketball specific endurance circuit training on aerobic capacity and heart rate of high school male basketball players. To achieve the purpose of the study twenty four (24) male high school basketball players were selected from Neyveli Lignite Corporation Sports School, Neyveli and St. Joseph Higher Secondary School, Manjakuppam, Cuddalore. These subjects were randomly distributed into two groups namely basketball specific endurance circuit training group (N=12) and control group (N=12). The mean age of the selected players was 16.85 ± 0.67 . Aerobic capacity, resting heart rate and peak heart rate were selected as criterion variables. Aerobic capacity was measured by multistage fitness test and resting and peak heart rate was measured using polar heart rate monitor. The basketball specific endurance circuit training was administered 3 days per week for six week. They performed 2 minutes of work at 90 to 95% of targeted heart rate using Karvonen method. They performed 8 repetitions during first and second week, followed by 10 repetitions during third and fourth week and 12 repetitions during fifth and sixth week of training. This was followed by 2 minutes of active resting at 60 to 70% of targeted heart rate. In this study 1:1 work rest ratio was followed. Both the groups were tested before and after training, the collected data was analysed using ANCOVA. The result of the study showed that aerobic capacity, resting heart rate and peak heart rate between the groups was significant, it indicate that after adjusting pre-test scores, there was a significant difference between the two groups on post-test scores. The findings of the study show that significant increase in aerobic capacity and decrease in resting and peak heart rate. It can be concluded that basketball specific endurance circuit training is effective in improving aerobic capacity and increases the cardiovascular fitness of male high school boys during competitive phase.

Keywords: *Endurance, Circuit training, resting heart rate, peak heart rate, aerobic capacity.*

Introduction

Traditionally, the coaches and trainers have planned conditioning programs for their teams by following regimens used by teams that have successful win-loss records. This type of reasoning is not sound because win-loss records alone do not scientifically validate the conditioning programs used by the successful teams. In fact, the successful team might be victorious by virtue of its superior athletes and not its outstanding conditioning program. Without question, the planning of an effective athletic conditioning program can best be achieved by the application of proven physiological training principles. Optimizing training programs for athletes is important because failure to properly condition an athletic team results in a poor performance and often defeat.

The importance of developing good conditioning programs based on the specific physiological demands of each sport is considered a key factor to success [1-3]. The basketball player needs to train multiple components of fitness. Thus, the athlete will concurrently perform various modes of training (e.g., strength, anaerobic, endurance). In the present study sport specific circuit training was employed. This incorporates skills and movements specific to the sport, at intensities sufficient to promote aerobic adaptations, are being increasingly implemented in professional team sports environment [4]. The perceived benefit of performing sports-specific exercise is that the training will transfer better into the athletes competitive environment and that the greatest training benefits occur when the training stimulus simulates the specific movement patterns and physiological demands of the sport [5]. The purpose of the study was to evaluate the effectiveness of a basketball specific endurance circuit training on aerobic capacity and heart rate of high school male basketball players.

Methods

Subjects

A total of twenty four (24) male high school basketball players were selected from Neyveli Lignite Corporation Sports School, Neyveli and St. Joseph Higher Secondary School, Manjakuppam, Cuddalore. These subjects were randomly distributed into two groups namely basketball specific endurance circuit training group (N=12) and control group (N=12). The mean age of the selected players was 16.85 ± 0.67 . The selected players had 3.8 ± 3.1 years of playing experience and regularly participate in training prior to the commencement of this study. All subjects were subjected to medical examination by a general medical

practitioner before participation in the study to ensure that there was of sufficient standard to be able to take part in fitness testing and training.

Variables and tests

Aerobic capacity, resting heart rate and peak heart rate were selected as criterion variables. Aerobic capacity was measured by multistage fitness test and resting and peak heart rate was measured using polar heart rate monitor.

Design of the study

For the present study pretest – posttest randomized group design [6] which consists of a control group (CG) and an experimental group (TG) that was used to find out effect sports specific circuit training on the selected physiological variables. Equal numbers (twelve) of subjects were assigned randomly to all the groups. TG was exposed to training with a set of drills selected for specific purpose. The TG underwent training for a period of six weeks (42 days). The training sessions were conducted three days a week (i.e. Monday, Wednesday, and Friday). Measurement of physiological variables was taken for both the groups.

Collection of Data

All the subjects were tested on physiological variables prior to training and after six weeks of training at Neyveli and Cuddalore. The testing session consists of warm-up and test interspersed with rest. All tests were explained and demonstrated. Before testing, subjects were given practice trials to become familiar with the testing procedures. All tests were counterbalanced pre and post testing to ensure that testing effects were minimized. Subjects performed each test as per test procedure and the scores of best trials were taken for this study. In the morning of the first day of testing measurements like height, weight, body composition, resting heart rate, vertical jump and repeated sprint ability were measured, however in the evening aerobic capacity and peak heart rate were evaluated.

Sports specific circuit training

TG is supplemented with sports specific circuit training replaced the regular physical fitness activity. However, control group performed regular physical activity. The training was carried out in outdoor basketball court. This sports specific circuit training was based on a previous design [7] and adapted to mimic as closely as possible the movement patterns of basketball match play as reported by McInnes *et al.*, (1995) [8]. The sports specific circuit training was administered 3 days per week for six week. The TG performed 2minutes of work at 90 to 95% of targeted heart rate using Karvonen method. They performed 8 repetitions during first and second week, followed by 10 repetitions during third and fourth week and 12 repetitions during fifth and sixth week of training. This was followed by 2 minutes of active resting at 70 to 80% of targeted heart rate. In this study 1:1 work rest ratio was followed. This training protocol was adapted from Helgerud *et al.*, (2001) [9]. The average running time of one circuit was 59 s and the total distance covered during one lap was approximately 153 m, with 60.2% of the movements forward sprinting and 39.8% side shuffling. The portion of the circuit considered „offence“ activity where a basketball was dribbled, was 55.6% while 44.4% was considered „defensive“ activity without ball. Three layups, three rebounds, seven vertical jumps, one pivot and 20 change of direction were completed during one repeat of the circuit.

Heart rate monitor was used to measure peak heart rate when performing the circuit. The subjects wore polar heart rate transmitter belt and watch (Polar heart rate monitor watch, Finland). The training intensity was fixed between 90 to 95% of THR. When the players perform below or above the prescribed intensity the watch will produce beep sound to alter their intensity accordingly. The sports specific endurance circuit training details are presented in figure 1.




Figure 1: Basketball Specific Circuit Training

The description of the circuit:

1-2 forward sprint; 2-3 hurdle jump; 3-4 forward sprint; 4 pivot left; 4-5 shuffle left; 5-6 shuffle right; 6-7 shuffle left; 7-8 shuffle right; 8-9 shuffle left; 9-10 shuffle right; 10-11 hurdle jump; 12 vertical jump (collect ball upon landing); 13-14 Zig Zag Dribble; 14-15 speed dribble with complete layup; 15 collect the rebound; 15-16-15 speed dribble with complete layup; 15 collect the rebound; 15-17-15 speed dribble with complete layup; 15 collect the rebound; 15-18 run and place the ball in basket; 18 throw the medicine ball; 18-19-20 forward sprint.

Statistical technique

The collected data was evaluated using Analysis of Covariance (ANCOVA). The proposed hypothesis was tested at 0.05 level of confidence. Beside this mean and standard deviation were also calculated. SPSS statistic software package (SPSS Company, America, version 17.0) was used. The α value of 0.05 was set for statistical significance.

Results

Table 1 clearly shows that aerobic capacity, resting heart rate and peak heart rate between the groups was significant, it indicate that after adjusting pre-test scores, there was a significant difference between the two groups on post-test scores on aerobic capacity, resting heart rate and peak heart rate. The findings of the study show that significant increase in aerobic capacity and decrease in resting and peak heart rate. The changes are presented in table 1.

Table 1
Changes in aerobic capacity and heart rate

Variables	Groups	Pre-test	Post-test	% of changes	F
Aerobic Capacity (ml/kg/min)	TG	43.36 \pm 5.82	45.12 \pm 5.42	3.29	7.890* ($p = 0.011$)
	CG	43.31 \pm 3.68	43.86 \pm 3.60	1.03	
Resting HR (beats/min)	TG	55.66 \pm 2.53	52.08 \pm 2.23	6.43	28.05* ($p = 0.000$)
	CG	54.50 \pm 2.57	54.75 \pm 2.83	0.42	
Peak HR (beats/min)	TG	198.58 \pm 3.57	192.33 \pm 3.82	3.14	32.20* ($p = 0.000$)
	CG	197.08 \pm 4.81	197.67 \pm 3.22	0.29	

Discussion

In the present study, basketball specific endurance circuit training for six week has significantly improved aerobic capacity 3.29%. Similarly, in CG 1.03% of improvement is elicited in aerobic capacity. The changes observed in the present study have been reported previously in basketball [9, 10]. The changes elicited in the present study found to be lower than the 7.5 to 9% increases in VO_{2peak} observed in soccer players following eight to ten-weeks of performing a similar sport-specific aerobic endurance training circuit compared to control group [11-12].

The reasons for small change obtained in aerobic capacity was firstly, differences observed could be due to the fact that the training was carried out during the competitive phase in the present study compared to the preparatory phase in other studies. Greater training adaptations are more likely to occur due to a potentially detrained state during preparatory phase. Secondly, the difference could also be due to the shorter duration training programme in the present study compared to others. Sports specific endurance circuit training results in increase capillary and mitochondrial density, enzyme activity (creatine phosphokinase and myokinase), metabolic stores (ATP, Creatine phosphate and glycogen), connective tissue strength (ligament and tendon) [13, 14]. These factors result in slight improvement in aerobic capacity in male high school basketball players.

Resting heart rate refers to the number of times a heart contracts in one minute (beats per minute or BPM) while at complete rest. The normal heart rate depends upon your age, gender and health and can vary greatly for both athletes and non-athletes. In general, a person's resting heart rate indicates their basic fitness level. The stronger the heart, the more blood it can pump during each contraction, and the less frequently it needs to beat to get adequate blood flow (circulation) and oxygen to the body tissues. A well trained athlete can have a very low resting heart rate and pump more blood than an unconditioned individual. In the present study TG showed 3.58 beats/min changes is elicited. The percentage reduction for resting heart rate

between pre to post was 6.48%. These changes are elicited as a result of sports specific endurance circuit training imparted to the high school male basketball players. The amount of blood pumped out of the left ventricle of heart with each contraction is called the stroke volume. Although some conditions can affect a person's stroke volume, endurance and high intensity cardiovascular exercise training often increases stroke volume [15]. A larger stroke volume results in a lower (resting) heart rate [16]. However, longer diastole influences the resting heart rate in athletes.

In this study TG showed 3.14% (6.25 beats/min) of reductions in peak heart rate. These alterations are caused because of sports specific endurance circuit training which resulted in improvement of aerobic capacity. Heart rate increases in parallel with increasing exercise intensity. Heart rate is stimulated to increase through the activation of mechano-, chemo- and baroreceptors sending afferent signals to the cardiovascular control centre in the brain. This in turn adjusts sympathovagal balance to the SA node bringing about a change in HR. At the onset of exercise, there is a rapid increase in HR. Due to its speed of response, this is suggested to arise through a withdrawal of parasympathetic modulation which enables the HR to increase up to the intrinsic rate of approximately 100 beats/min. Thereafter, any increase in HR is stimulated through an increased sympathetic modulation. Increased sympathetic cardiac modulation is evident from approximately 25% peak VO_2 onwards and by the time exercise reaches an intensity of 50–60% of peak VO_2 , data suggest that vagal modulation disappears all together. Very few studies have reported the dynamics of autonomic control of HR during exercise in children. Those studies that have been performed report similar findings to those observed in adults. Due to training adaptations these changes are found in the present study.

Conclusion

Basketball specific endurance circuit training is effective in improving aerobic capacity and increases the cardiovascular fitness of male high school boys during competitive phase.

References

- [1] G.M. Gillam, Physiological basis of basketball bioenergetics, *NSCA Journal*, 6 (1984) 44-71.
- [2] J. Taylor, Basketball: applying time motion data to conditioning, *Strength and Conditioning Journal*, 25(2003) 57-64.
- [3] J. Taylor, A tactical metabolic training model for collegiate basketball, *Strength and Conditioning Journal*, 5 (2004) 22-29.
- [4] E. Lawson, Incorporating sports-specific drills into conditioning. In B. Foran (Ed.), *High performance sports conditioning* (2001) (pp. 215-266). Champaign, IL: *Human Kinetics*.
- [5] D.M. McArdle, F.I. Katch, V.L. Katch, Exercise physiology: energy, nutrition and human performance (5th Ed.). (1996) Philadelphia, PA: Lippincott Williams and Wilkins.
- [6] J.R. Thomas, J.K. Nelson, and S.J. Silverman, Research Methods in Physical Activity, (1996) USA.
- [7] M. Smith, Basketball skill test for the big men, *FIBA Assist Magazine*, 07 (2004) 59-60.
- [8] S.E. McInnes, J.S. Carlson, C.J. Jones, M.J. McKenna, The physiological load imposed on basketball players during competition, *Journal of Sports Science*, 13 (1995) 387-97.
- [9] J. Helgerud, L.C. Engen, U. Wisloff, J. Hoff, Aerobic endurance training improves soccer performance, *Medicine & Science in Sports & Exercise*, 33 (2001) 1925-31.
- [10] C.P. Balabinis, C.H. Psarakis, M. Moukas, M.P. Vassiliou, P.K. Behrakis, Early phase changes by concurrent endurance and strength training, *Journal of Strength and Conditioning Research*, 17 (2003) 393-401.
- [11] K. Chamari, Y. Hachana, F. Kaouech, R. Jeddi, I. Moussa-Chamari, U. Wisloff, Endurance training and testing with the ball in young elite soccer players, *British Journal of Sports Medicine*, 39 (2005) 24-8.
- [12] K. McMillan J. Helgerud, R. Macdonald, and J. Hoff, Physiological adaptations to soccer specific endurance training in professional youth soccer players, *British Journal of Sports Medicine*, 39 (2005) 273-277.
- [13] T.R. Baechle, and R.W. Earle, NSCA Essentials of strength training and conditioning (2nd ed.) (2000) *Human Kinetics*, Leeds, UK.
- [14] N. Amigo, J.A. Cadefau, I. Ferrer, N. Tarados, and R. Cusso, Effect of summer intermission on skeletal muscle of adolescent soccer players, *Journal of Sports Medicine and Physical Fitness*, 38 (1998) 298–304.
- [15] D. Bonaduce, M. Petretta, V. Cavallaro, C. Apicella, A. Ianniciello, M. Romano, R. Breglio, F. Marciano, Intensive training and cardiac autonomic control in high level athletes, *Medicine & Science in Sports & Exercise*, 30 (1998) 691-6.
